

SUMMIT DIGESTIVE & LIVER DISEASE SPECIALISTS

**DAVID CHUA, MD
RAJEEV NAYAR, MD
ALBERT SAPORTA, MD**

BOWEL PREPARATION FOR CLENPIQ

FIVE DAYS BEFORE YOUR COLONOSCOPY:

- Aim to have a light, low fiber diet (50% less of what you regularly eat) to reduce stool production
- Avoid nuts, lettuce, fruits and vegetables with seeds and popcorn
- Take Milk of Magnesia 3 tablespoons every night at bedtime
- Stop taking iron supplements and NSAIDs (Ibuprofen, Advil, Motrin, Aleve, Naproxen)

THE DAY BEFORE YOUR PROCEDURE: Follow a clear liquid diet ALL day long (NO SOLID FOOD):

- Drink 8 oz (1 cup) of clear liquids every hour while awake
- Water (Plain, carbonated or flavored)
- Fruit juice without pulp (Apple juice or white grape juice)
- Lemonade or clear colored soda (Sprite, 7-up)
- Black coffee or tea (DO NOT ADD ANY MILK OR CREAM)
- Sports drinks such as Gatorade, Powerade, etc (NO RED OR PURPLE)
- Clear, fat-free broth (Chicken, beef or vegetable)
- Popsicles without ANY milk, fruit bits, seeds, nuts (NO RED OR PURPLE)

Clenpiq Instructions: **PLEASE FOLLOW THE INSTRUCTIONS BELOW. DO NOT FOLLOW THE INSTRUCTIONS ON THE BOX**

- 1:00PM the day before your procedure: Drink 1 bottle of Clenpiq followed by 5 8oz cups of any clear liquid.
- 6:00PM the day before your procedure: Take 4 tablets of Dulcolax followed by 2 8oz cups of any clear liquid.
- 9:00PM the day before your procedure: Drink 1 bottle of Clenpiq followed by 4 8oz cups of any clear liquid.

****IMPORTANT: YOU NEED TO STOP DRINKING LIQUIDS 6 HOURS BEFORE YOUR PROCEDURE TIME!****

****IMPORTANT: ASK A FRIEND OR RELATIVE TO DRIVE AND ASSIST YOU HOME AFTER THE PROCEDURE. YOU CANNOT TAKE AN UBER OR TAXI HOME ALONE!****

DIABETIC PATIENTS:

- If you are taking an oral hypoglycemic: **DO NOT** take it the day before and of your procedure.
- If you are taking **insulin**, please take half of your prescribed AM insulin dose the day before the procedure and **skip your night time insulin dose**.

MEDICATION INSTRUCTIONS:

- **You will need to stop the medications listed below 5 days before the procedure:**
Anticoagulants/blood thinners: Coumadin/Warfarin, Apixaban/Eliquis, Plavix/Clopidogrel, Rivaroxaban/Xarelto, Ticagrelor/Brilinta, Prasugrel/Effient.
Consult your prescribing doctor before stopping this medication.
- **You may take your regular medication unless instructed otherwise.**

If you have any questions please contact our office at 630-889-9889